arXiv:0905.1936v1 [astro-ph.EP] 12 May 2009

New astrometry and photometry for the companion candidates of CT Cha

Tobias O. B. Schmidt*, Ralph Neuhäuser†, Markus Mugrauer†, Ana Bedalov**,† and Nikolaus Vogt‡,§

*Astrophysikalisches Institut, Universität Jena, Schillergässchen 2-3, 07745 Jena, Germany, tobi@astro.uni-jena.de
†Astrophysikalisches Institut, Universität Jena, Schillergässchen 2-3, 07745 Jena, Germany
**Faculty of Natural Sciences, University of Split, Teslina 12. 21000 Split, Croatia
‡Departamento de Física y Astronomía, Universidad de Valparaíso, Avenida Gran Bretaña 1111, Valparaíso, Chile
§Instituto de Astronomía, Universidad Catolica del Norte, Avda. Angamos 0610, Antofagasta, Chile

Abstract. In our ongoing search for close and faint companions around T Tauri stars in the Chamaeleon star-forming region, we recently (Schmidt et al. 2008b) presented direct observations and integral field spectroscopy of a new common proper motion companion to the young T-Tauri star and Chamaeleon member CT Cha and discussed its properties in comparison to other young, low-mass objects and to synthetic model spectra from different origins. We now obtained for the first time direct H-Band observations of the companion CT Cha b and of another faint companion candidate (cc2) approximately 1.9 arcsec northeast of CT Cha using the Adaptive Optics (AO) instrument Naos-Conica (NACO) at the Very Large Telescope (VLT) of the European Southern Observatory (ESO) in February 2008. From these data we can now exclude by 4.4 & 4.8 σ that CT Cha b is a non-moving background object and find cc2 to be most likely a background star of spectral type $\leq$ K4 with a proper motion of $\mu_\alpha \cos\delta$= -8.5 $\pm$ 5.7 mas/yr and μ_δ= 12.0 $\pm$ 5.6 mas/yr, not consistent with being a member of the Cha I star-forming region.

Keywords: Stars: low-mass, brown dwarfs – Stars: pre-main sequence – Stars: planetary systems: formation – Stars: individual: CT Cha
PACS: 95.85.Jq, 97.21.+a, 97.82.Fs

INTRODUCTION

CT Cha (aka HM 9), introduced in the 65th Name-List of Variable stars by Kholopov et al. (1981), was originally found by Henize & Mendoza (1973) and later classified to be a classical T Tauri star by Weintraub (1990) and Gauvin & Strom (1992) from IRAS data (see Schmidt et al. (2008b) and references therein for more information on CT Cha A).

We recently (Schmidt et al. 2008a & 2008b), identified a very faint co-moving substellar companion, just ~ 2.67 arcsec northwest of CT Cha corresponding to a projected separation of ~ 440 AU as well as another companion candidate (cc2) ~ 1.9 arcsec northeast of the primary star. We could show that CT Cha A and CT Cha b form a common proper motion pair and that the companion is by $\geq 4\,\sigma$ significance not a stationary background object. Our near-infrared spectroscopy yielded a temperature of 2600 ± 250 K for the companion and an optical extinction of A_V= 5.2 ± 0.8 mag,

TABLE 1. VLT/NACO observation log. Remark: Each image consists of the number of exposures given in column 4 times the individual integration time given in column 3.

JD - 2453700 [days]	Date of observation	DIT [s]	NDIT	No. of images	Filter
83.54702	17 Feb 2006	1.5	25	20	Ks
460.62535	1 Mar 2007	4	15	21	J
461.63001	2 Mar 2007	4	7	30	Ks
815.81036	19 Feb 2008	1	60	5	H

TABLE 2. Absolute and relative astrometric results for CT Cha A and CT Cha b. (a) PA is measured from N over E to S. (b) Relative changes are given relative to epoch 3 at JD 2454515.81036.

JD - 2453700 [days]	Pixel scale [mas/Pixel]	Separation [arcsec]	PA^a [deg]	Change in separation [pixel]	Change in PA^a [deg]
83.54702	13.24 ± 0.18	2.670 ± 0.036	300.71 ± 1.24	-0.19 ± 0.36	0.219 ± 0.166
461.63001	13.24 ± 0.19	2.670 ± 0.038	300.68 ± 1.32	-0.28 ± 0.18	0.183 ± 0.082
815.81036	13.24 ± 0.20	2.674 ± 0.040	300.50 ± 1.40	^{b}	^{b}

when compared to spectra calculated from Drift-Phoenix model atmospheres (Helling et al. 2008). We further demonstrated the validity of the model fits by comparison to several other well-known young sub-stellar objects. Due to a prominent Pa-β emission in the J-band, accretion is probably still ongoing onto CT Cha b. From temperature and luminosity ($\log(L_{bol}/L_{\odot}) = -2.68 \pm 0.21$), we derived a radius of $R = 2.20^{+0.81}_{-0.60}\,R_{Jup}$. We found a consistent mass of $M = 17 \pm 6\,M_{Jup}$ for CT Cha b from both its luminosity and temperature when placed on evolutionary tracks. Hence, CT Cha b is most likely a wide brown dwarf companion or possibly even a planetary mass object, depending on both the true mass of CT Cha b and the definition of planets.

ASTROMETRY

CT Cha b

We observed CT Cha in three epochs in February 2006, March 2007 and in February 2008 (see Table 1). All observations were done with the European Southern Observatory (ESO) Very Large Telescope (VLT) instrument Naos-Conica (NACO).

To check again for the common proper motion of CT Cha b we have repeated the analysis done in Schmidt et al. (2008b) and included the new H-Band epoch listed as last entry in Table 1. From Fig. 1 (left), we can now exclude by 4.4 & 4.8 σ that CT Cha b is a non-moving background object.

Companion Candidate 2 (cc2)

To check for the common proper motion of the tentative second companion (cc2) of CT Cha, we also repeated the analysis done in Schmidt et al. (2008b) for CT Cha



FIGURE 1. Observed separation between CT Cha A and CT Cha b (*left*) and cc2 (*right*). Our measurements from 2006, 2007, 2008 are shown. The long dashed lines enclose the area for constant separation, as expected for a co-moving object. The dash-dotted line is the change expected if CT Cha b / cc2 is a non-moving background star. The opening cone enclosed by the continuous lines its estimated errors. The waves of this cone show the differential parallactic motion that has to be taken into account if the other component is a non-moving background star with negligible parallax. The opening short-dashed cone is for the combination of co-motion and the maximum possible orbital motion for a circular edge-on orbit.

TABLE 3. Apparent magnitudes of CT Cha b and CT Cha cc2. See Table 1 for JDs of the observations.

		CT Cha b			CT Cha cc2	
Epoch	J-band	H-band	Ks-band	J-band	H-band	Ks-band
2006			14.95 ± 0.30			17.36 ± 0.31
2007	16.61 ± 0.30		14.89 ± 0.30	18.15 ± 0.32		17.27 ± 0.31
2008		15.56 ± 0.30			17.48 ± 0.32	

b for this object and included the new H-Band epoch (see Table 1). As can be seen in Fig. 1, the companion candidate cc2 is moving like a stationary background object in separation from CT Cha A and, hence, is not a low-mass companion. Given the change in separation and position angle, the proper motion of CT Cha cc2 $\mu_\alpha \cos\delta$= -8.5 ± 5.7 mas/yr and μ_δ= 12.0 ± 5.6 mas/yr is by $\sim 3\,\sigma$ not consistent with the median proper motion of 61 members of the Cha I star-forming region of $\mu_\alpha \cos\delta$= -21 ± 1 mas/yr and μ_δ= 2.0 ± 1 mas/yr (Luhman et al. 2008).

PHOTOMETRY

In addition we compute from our NACO measurements also H-band photometry from the flux ratio of the companion candidates relative to CT Cha A (see Table 3). From the J-, H- and Ks-band magnitudes we can now present an artificial color-composite image, shown in Fig. 2. If we assume the same amount of extinction for cc2 as we found for CT Cha A, i.e. $A_V \sim 1.3$ mag (Schmidt et al. 2008b) or possibly more, as the star might be a background star, we find a spectral type of $\leq$ K4 from the J-, H-, and K-band photometry



FIGURE 2. Artificial color-composite image from our J-, H- and Ks-band images taken in 2007 and 2008. CT Cha b was found $\sim$2.7 arcsec northwest (marked) of the classical T Tauri star CT Cha A. Northeast of CT Cha A the background star cc2 has currently a separation of $\sim$1.9 arcsec.

given in Table 3 in comparison to the colors of main-sequence stars given in Kenyon & Hartmann (1995), hence, at the main-sequence, a distance of ≥ 350 pc.

ACKNOWLEDGMENTS

TOBS would like to thank Evangelisches Studienwerk e.V. Villigst for financial support. NV acknowledges support by FONDECYT (grant 1061199) and Universidad de Valparaiso (grant 07/2007). AB and RN wish to acknowledge support from the German National Science Foundation (Deutsche Forschungsgemeinschaft, DFG), grants NE 515/13-1 and 13-2.

REFERENCES

1. L. S. Gauvin, and K. M. Strom, *ApJ* **385**, 217–231 (1992).
2. Ch. Helling, M. Dehn, P. Woitke, and P. H. Hauschildt, *ApJ* **675**, L105–L108 (2008), `arXiv:0801.3733`.
3. K. G. Henize, and E. E. Mendoza, *ApJ* **180**, 115–119 (1973).
4. S. J. Kenyon, and L. Hartmann, *ApJS* **101**, 117–+ (1995).
5. P. N. Kholopov, N. N. Samus’, N. P. Kukarkina, G. I. Medvedeva, and N. B. Perova, *Informational Bulletin on Variable Stars* **1921**, 1–+ (1981).
6. K. L. Luhman, L. E. Allen, P. R. Allen, R. A. Gutermuth, L. Hartmann, E. E. Mamajek, S. T. Megeath, P. C. Myers and G. G. Fazio, *ApJ* **675**, 1375–1406 (2008).
7. T. Schmidt, and R. Neuhäuser, “Finding new sub-stellar co-moving companion candidates - the case of CT Cha,” in *IAU Symposium*, 2008a, vol. 249 of *IAU Symposium*, pp. 65–69.
8. T. O. B. Schmidt, R. Neuhäuser, A. Seifahrt, N. Vogt, A. Bedalov, C. Helling, S. Witte, and P. H. Hauschildt, *ArXiv e-prints* **809** (2008b), `0809.2812`.
9. D. A. Weintraub, *ApJS* **74**, 575–594 (1990).